

BOTRY-Zen®WP

Botrytis, Sclerotinia

Mix 8g (approx. half a bottle) per litre of water. Pre-mix powder with a small amount of water first before adding in remaining water. Ensure powder is mixed thoroughly. Apply early and late bloom. Further applications can be made up to 5 days before picking. Ensure new growth at risk is protected by applying early. Keep pressure spray bottle well shaken throughout application.

PRECAUTIONS

Store in original container, tightly closed, away from foodstuffs. Avoid inhalation of product dust or spray mist. Wear a face mask when handling BOTRY-Zen®WP. Dispose of unused or expired material by dilution with water and disposal to headlands. In case of spillage, wash area with 10% bleach. Not environmentally hazardous

STORAGE

Store in a cool dry place (below 20°C). Keep away from light

SHELF-LIFE

This product has a 1 year shelf life.

MIDI-Zen®

Botrytis, Powdery Mildew

Mix 30ml (approx. one bottle) per litre of water. Shake MIDI-Zen® first before mixing with water. Ensure it is mixed thoroughly. Apply at first signs of infection. Continue applications every 10-14 days if infection appears. Keep pressure spray bottle well shaken throughout application.

Hazard Class Ecotoxic 9.1D Designed for biocidal action. Avoid contamination of all waterways. It is an offense to use this product on animals.

PRECAUTIONS

Do not exceed the recommended application rate as this can cause phytotoxicity. Store in original bottle tightly closed, away from foodstuffs. Dispose of unused or expired material by dilution with water. In case of spillage, wash area.

STORAGE

Store in original packaging. Keep away from light and store at <20°C.

COMPATIBILITY

MIDI-Zen is compatible with BOTRY-Zen and Topas. Do not add other pesticides unless they have been specifically cleared as being compatible

SHELF-LIFE

This product has a 2 year shelf life.

AMOUR-Zen®

Botrytis, Sclerotinia, Downy Mildew

Mix 15ml (approx. half a bottle) per litre of water. Ensure it is mixed thoroughly. Apply before periods of disease and then 7 – 10 days later. Continue applications every 7 – 10 days if infection appears. Keep pressure spray bottle well shaken throughout application.

PRECAUTIONS

Keep out of reach of children. Store in original container tightly closed away from foodstuffs. Avoid contact with skin and eyes and inhalation of spray mist and/or concentrate. When mixing or applying, wear protective eyewear and waterproof gloves. Wash hands and exposed skin before meals and after using. This substance must not be applied onto or into water (other than for the purpose of spraying)

COMPATIBILITY

Do not mix ARMOUR-Zen® with other biological fungicides. When using ARMOUR-Zen® do not apply calcium solutions.

SHELF-LIFE

This product has a 4 year shelf life

BOTRYZEN NATURAL GARDEN DISEASE CONTROL™



**USE PRESSURED
SPRAY BOTTLE
ONLY. SPRAY
THOROUGHLY
UNTIL SPRAY
RUNS OFF THE
LEAVES.**

FIRST AID

If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or call a doctor immediately. If skin contact occurs, wash off thoroughly with disinfectant. If splashed in eyes, wash out immediately with water. Seek medical attention.

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS

Registered to:
Botry-Zen (2010) Limited
21 Willis Street
Dunedin, New Zealand
info@botryzen.co.nz
www.botryzen.co.nz
Phone: 03 477 6447

See www.nzfsa.govt.nz/acvm/ for registration conditions